



**PRESENTATION:** Film-Coated tab 100 mg x 1 x 10's. Oral Susp 40 mg/5 mL x 50 ml.

**ACTIONS:** Cefpodoxime proxetil is an orally administered, extended spectrum, semisynthetic antibiotic of the third generation.

**Microbiology:** Cefpodoxime is active in vitro against a wide range of gram-positive and gram-negative bacteria.

**Note:** Cefpodoxime is inactive against most strains of Enterococcus, Pseudomonas and Enterobacter.

**INDICATIONS:** Treatment of patients with mild to moderate infections caused by susceptible strains of the following organisms:

**Skin and Skin Structure:** Uncomplicated skin and skin structure infections caused by Staphylococcus aureus.

**Upper Respiratory Tract Infections:** Acute otitis media caused by Streptococcus pneumoniae, Haemophilus influenzae.

**DOSAGE & ADMINISTRATION:** Film-Coated Tablet Cefprox tablet should be administered orally with food.

**Renal impairment:** For patients with severe renal impairment (creatinine clearance <30 ml/min), the dosage should be reduced.

**OVERDOSAGE:** In the event of serious toxic reaction from overdosage, haemodialysis or peritoneal dialysis may be beneficial.

**CONTRAINDICATIONS:** Known allergy to cefpodoxime or to any of the antibiotics in the cephalosporin class.

**WARNINGS:** Before therapy with Cefprox is instituted, careful inquiry should be made to determine whether the patient has a history of allergic reactions to any of the antibiotics in the cephalosporin class.

**PRECAUTIONS:** **Carcinogenicity:** Long-term animal carcinogenesis studies of cefpodoxime proxetil have not been conducted.

**Mutagenicity:** Mutagenesis studies of cefpodoxime were all negative.

**Use in pregnancy:** There are no adequate and well-controlled studies of cefpodoxime proxetil use in pregnant women.

**Labor and Delivery:** Cefprox has not been studied for use during labor and delivery. Treatment should not be initiated during labor.

**Use in lactation:** Cefpodoxime is excreted in human milk. Because of the potential for serious reactions in nursing infants, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

**Use in children:** Safety and efficacy in infants under 5 months have not been established.

**ADVERSE REACTIONS:** Film-Coated Tablets (Multiple Dose): In clinical trials using multiple doses of cefpodoxime proxetil, the following adverse reactions were observed:

**Dermatologic:** Fungal skin infection, skin scaling/peeling. **Endocrine:** Menstrual irregularity.

**Genital:** Pruritus.

**Gastrointestinal:** Flatulence, decreased salivation, candidiasis, pseudomembranous colitis.

**Hypersensitivity:** Anaphylactic shock.

**Metabolic:** Decreased appetite.

**Miscellaneous:** Malaise. Fever.

**Central Nervous System:** Dizziness, fatigue, anxiety, insomnia, flushing, nightmares, weakness.

**Respiratory:** Cough epistaxis. **Special Senses:** Altered taste, eye itching, tinnitus.

**Oral Suspension (Multiple Dose):** In clinical trials using multiple doses of cefpodoxime proxetil granules, the following adverse reactions were observed:

**Incidence <1%:** **Central Nervous System:** Headache, irritability.

**Dermatologic:** Exacerbation of acne, exfoliative dermatitis.

Genital: Pruritus or vaginitis.

Gastrointestinal: Nausea, abdominal pain, candidiasis, decreased salivation, pseudomembranous colitis.

Metabolic: Decreased appetite.

Miscellaneous: Fever.

Psychiatric: Hyperactivity/nervousness.

Respiratory: Epistaxis, rhinitis.

Laboratory Changes: Significant laboratory changes that have been reported in clinical trials of cefpodoxime proxetil include:

Haematologic: Eosinophilia, leukocytosis, lymphocytosis, granulocytosis, basophilia, monocytosis, thrombocytopenia.

Serum Chemistry: Increased glucose, decreased glucose, decreased serum albumin, decreased serum calcium, decreased serum potassium.

Renal: Increased BUN and creatinine levels. Most of these abnormalities were transient and not clinically significant.

INTERACTIONS: Antacids: Concomitant administration high doses of antacids (sodium bicarbonate and sodium citrate) may decrease the absorption of cefpodoxime proxetil.

Probenecid: As with other  $\beta\beta$ -lactam antibiotics, renal excretion of cefpodoxime was inhibited by probenecid.

Drug/Laboratory Test Interactions: None known. Other drugs including cefpodoxime proxetil are known to occasionally interact with laboratory tests.

Table 1.			
Adults (age $\geq$ 13 years)	Total Daily Dose	Dose Frequency	Duration
Acute community acquired pneumonia	400 mg	200 mg every 12 hrs	14 days
Acute bacterial exacerbations of chronic bronchitis	400 mg	200 mg every 12 hrs	10 days
Uncomplicated gonorrhoea and rectal gonococcal infections	200 mg	Single dose	7-14 days
Sten and skin structure	800 mg	400 mg every 12 hrs	5-10 days
Pharyngitis and/or tonsillitis	200 mg	100 mg every 12 hrs	7 days
Uncomplicated urinary tract infection	200 mg	100 mg every 12 hrs	7 days

  

Table 2.			
Children (age 5 months through 12 years)	Total Daily Dose	Dose Frequency	Duration
Acute otitis media	8 mg/kg/day (Max 400 mg/day)	4 mg/kg every 12 hrs (Max 200 mg/dose)	5 days
Pharyngitis and/or tonsillitis	8 mg/kg/day (Max 200 mg/day)	4 mg/kg every 12 hrs (Max 100 mg/dose)	5-10 days

Любой парнишка может сломать ногу.

Рэндалл, вы "[Все альбомы three days grace скачать](#)" думающий человек, вы работаете больше многих "

[Neverwinter nights 2 патчи](#)

"других, и вам удастся сохранять чувство юмора, когда речь заходит "

[Скачать караоке на компьютере с баллами](#)

"о происходящем.

Женщина-паук заберется на вашу "[Король и шут todd скачать](#)" скалу, Я поздравляю с вами, Но пойду другой дорогой.

Казалось, что "[Программы для lg gs290](#)" тут был еще кто-то.

Он повернул вправо и направился к полю, "[Игры alawar полные версии](#)" что находилось в трех четвертях мили.

document.getElementById("J#1367253187bl3ed0106c").style.display = "none";

Он не всегда делает это, во всяком случае, "[Скачать бунин лёгкое дыхание](#) "когда не чувствует себя как надо, объяснил мальчик.

Я опустил "[Бесплатная музыка скачать шуфутинский](#) "стилет во внутренний карман и повернулся.

Да, они выглядят как обычные трассеры, только "[Игры бесплатно для виндовс 7](#) "голубого цвета.

Я закрыл дверь, но эхо "[Мотоциклы игры для мальчиков бесплатно](#) "ударов ножей о мишень долго и неотступно преследовало меня.

Подобного празднества она не видела ни разу в жизни.

К этим терпким "[Скачать игру f1 2010](#) "ароматам порой примешивался легкий дух Европы.

document.getElementById("J#1368125624bl35be13c9").style.display = "none";